

# Distinguishing adaptive from nonadaptive genetic differentiation: comparison of $Q_{ST}$ and $F_{ST}$ at two spatial scales

S Volis<sup>1</sup>, B Yakubov<sup>2</sup>, I Shulgina<sup>3</sup>, D Ward<sup>4</sup> and S Mendlinger<sup>3</sup>

<sup>1</sup>Life Science Department, Ben-Gurion University of the Negev, POB 653, Beer Sheva 84105, Israel; <sup>2</sup>Horticulture Department, Purdue University, 170 S. University Avenue, West Lafayette, IN 47907, USA; <sup>3</sup>The Jacob Blaustein Institute for Desert Research, Ben-Gurion University of the Negev, Sede Boqer 84990, Israel; <sup>4</sup>School of Biological and Conservation Sciences, University of KwaZulu-Natal, P. Bag X01, Scottsville 3209, South Africa

Genetic differentiation in 20 hierarchically sampled populations of wild barley was analyzed with quantitative traits, allozymes and Random Amplified Polymorphic DNAs (RAPDs), and compared for three marker types at two hierarchical levels. Regional subdivision for both molecular markers was much lower than for quantitative traits. For both allozymes and RAPDs, most loci exhibited minor or no regional differentiation, and the relatively high overall estimates of the latter were due to several loci with exceptionally high regional differentiation. The allozyme- and RAPD-specific patterns of differentiation were concordant in general with one another, but not with quantitative trait differentiation. Divergent selection on quantitative traits inferred from very high regional  $Q_{ST}$  was in full agreement with our previous results obtained from a test of local adaptation and multilevel selection analysis. In contrast,

most variation in allozyme and RAPD variation was neutral, although several allozyme loci and RAPD markers were exceptional in their levels of regional differentiation. However, it is not possible to answer the question whether these exceptional loci are directly involved in the response to selection pressure or merely linked to the selected loci. The fact that  $Q_{ST}$  and  $F_{ST}$  did not differ at the population scale, that is, within regions, but differed at the regional scale, for which local adaptation has been previously shown, implies that comparison of the level of subdivision in quantitative traits, as compared with molecular markers, is indicative of adaptive population differentiation only when sampling is carried out at the appropriate scale.

*Heredity* advance online publication, 28 September 2005; doi:10.1038/sj.hdy.6800745

**Keywords:** adaptation; population genetic structure; molecular markers; *Hordeum spontaneum*

## Introduction

Evolutionary forces determining population subdivision and the extent of genetic variation include those influencing all loci equally (migration, genetic drift and mating system) and those affecting loci differentially (natural selection and mode of inheritance). Cavalli-Sforza (1966) introduced the idea that heterogeneity of population differentiation ( $F_{ST}$ ) values across loci reflects the involvement of different loci in an adaptive process. Subsequently, Lewontin and Krakauer (1973) further developed this idea into a formal statistical test. Until recently, allozyme loci encoding functional enzymes were the only molecular markers available to examine this idea, and, despite the continuing debate about possible direct selection effects on allozymes (Lewontin, 1991), allozymes have been assumed to be selectively neutral in most studies. The quantitative traits that represent additive or interactive effects of many contributing loci are more directly related to fitness than molecular markers; that is, while many may be neutral, some quantitative traits are adaptive. Selectively neutral

quantitative traits are predicted to behave similarly to traits determined by a single locus (Rogers and Harpending, 1983). This led to the idea of testing selective divergence in quantitative traits by comparison of their population differentiation ( $Q_{ST}$ ) with the population differentiation obtained from neutral molecular markers ( $F_{ST}$ ) (Spitze, 1993). Three outcomes, each having a unique interpretation, are possible (Merila and Crnokrak, 2001): (i)  $Q_{ST} > F_{ST}$ : this is commonly interpreted as evidence of divergent selection and adaptation to local environments (Podolsky and Holtsford, 1995; Bonnin *et al.*, 1996; Latta and Mitton, 1997; Jaramillo-Correa *et al.*, 2001; Steinger *et al.*, 2002; Storz, 2002; Chan and Arcese, 2003; Luttikhuisen *et al.*, 2003); (ii)  $Q_{ST}$  and  $F_{ST}$  do not differ: genetic drift alone is sufficient to explain the pattern of differentiation (Yang *et al.*, 1996), although effects of natural selection and drift in certain cases can be indistinguishable in certain cases (Sokal and Wartenburg, 1983); and (iii)  $Q_{ST} < F_{ST}$ : convergent selection favoring the same phenotype in different environments is assumed (Kuittinen *et al.*, 1997; Petit *et al.*, 2001; Edmands and Harrison, 2003). However, interpretation of any discrepancy between  $Q_{ST}$  and  $F_{ST}$  as evidence of selection on a particular quantitative trait may not always hold. As variation for quantitative traits is introduced by mutation at a higher rate than for

molecular markers (Kimura, 1983; Lynch, 1988), the extent of variation for quantitative traits usually exceeds that for molecular markers (Morgan *et al*, 2001). The population divergence due to stochastic processes such as drift and founder effects may be detectable with quantitative traits but not with molecular markers when the latter possess a low level of polymorphism or when the genes involved in the expression of quantitative traits have significant epistatic effects (Lynch, 1994). One way to clarify the issue is to conduct a test of local adaptation. Unfortunately, no rigorous test of local adaptation has been conducted in a comparative study of Q<sub>ST</sub> and F<sub>ST</sub>.

Different classes of molecular markers may also differ in the level of population differentiation they reveal because of differences in mutation rate, neutrality or linkage disequilibrium with other loci that are subject to selection. In several studies, lower G<sub>ST</sub> (between population diversity) values were found for allozymes than Random Amplified Polymorphic DNAs (RAPDs) (Ayres and Ryan, 1997; Latta and Mitton, 1997; Comes and Abbott, 2000), although other studies detected no difference (Isabel *et al*, 1995; Szmids *et al*, 1996; Le Corre *et al*, 1997; Aagaard *et al*, 1998) or lower population differentiation in RAPDs has occasionally been detected (Allendorf and Seeb, 2000; Bartish *et al*, 2000b). As with quantitative *vs* DNA/allozyme markers, discordant patterns of divergence in several molecular markers and loci are commonly interpreted as the differential effect of natural selection on those markers and loci (Karl and Avise, 1992; McDonald, 1994; Latta and Mitton, 1997), despite no direct evidence of the latter.

Not only is the degree of population differentiation measured by F<sub>ST</sub> or Q<sub>ST</sub> important for understanding population genetics but also the pattern of population differentiation and especially the concordance (or discordance) in patterns of differentiation revealed by different markers. In a recent study where desert and Mediterranean plants of *Hordeum spontaneum* were found to be adapted to their locations and differentiated in phenotypic traits (Volis *et al*, 2002b), no concordance between quantitative and molecular (allozyme) pairwise genetic distances was detected (Volis *et al*, 2002a). Several studies in which patterns of quantitative trait variation were strongly associated with climate or distinct habitats also demonstrated a lack of correspondence between molecular and quantitative trait variation patterns (Karhu *et al*, 1996; Knapp and Rice, 1998; McKay *et al*, 2001; Steinger *et al*, 2002). In other studies where genetic subdivisions at quantitative and molecular markers were compared in their relation to geographic subdivision but without tests of local adaptation/association with environmental parameters, or were compared without any reference to spatial location/environmental differences, the outcomes varied from a close match (Lagercrantz and Ryman, 1990; Kuittinen *et al*, 1997; Morgan *et al*, 2001) to complete disagreement (Podolsky and Holtsford, 1995; Black-Samuelsson *et al*, 1997; Heaton *et al*, 1999; Storz, 2002). When only allozyme and RAPD population differentiation patterns based on genetic distances were compared, the correspondence varied from low (Mamuris *et al*, 1999; Sun and Wong, 2001; Vandewoestijne and Baguette, 2002) to very high (Peakall *et al*, 1995; Lifante and Aguinalgalde, 1996; Jenczewski *et al*, 1999).

There is a need for a robust approach to verify the common interpretations of observed patterns of differentiation listed above, viz. Q<sub>ST</sub> > F<sub>ST</sub> as evidence of divergent selection and Q<sub>ST</sub> < F<sub>ST</sub> as evidence of convergent selection (Bossart and Pashley Prowell, 1998; Merila and Crnokrak, 2001). To date, a limited number of studies have undertaken the development of such an approach (Ross, 1997; Ross *et al*, 1997, 1999; McKay *et al*, 2001; Volis *et al*, 2002a, b).

Self-fertilization is a life history character reducing effective population size and frequency of recombination and limiting efficient migration to seed dispersal only. Thus, self-fertilization decreases within-population diversity and greatly increases the potential for population genetic structuring as compared with outcrossing (Charlesworth, 2003). Local selection will exacerbate the above effects, leading to large allele frequency differences, and high F<sub>ST</sub> values in a large proportion of marker loci (Charlesworth *et al*, 1997). As pointed out by Charlesworth (2003) this implies that it is particularly difficult to identify loci subject to selection in inbreeders. Therefore, a comparison of F<sub>ST</sub> and Q<sub>ST</sub> in a selfing species is a more conservative test for neutral selectivity than in an outcrossing species because only strong local selection will cause Q<sub>ST</sub> > F<sub>ST</sub> in a selfer, while drift alone or moderate selection will result in approximately equal and high values of Q<sub>ST</sub> and F<sub>ST</sub>. On the other hand, the level of population subdivision may differ at different geographical (environmental) scales, and sampling may or may not be relevant to the actual scale at which adaptive differentiation occurs. Therefore, a selfing species sampled at several hierarchical levels may provide insights into how much (if any) local selection contributes to population subdivision in Q<sub>ST</sub> and F<sub>ST</sub> at the level relevant to local adaptation.

The present study extends our previous studies of life history and population genetic differentiation in wild barley (*H. spontaneum* Koch), sampled using hierarchical design in four different environments (regions) representing distinct climatic zones and vegetation types. The 20 sampled populations (five populations per region) were analyzed for allozyme and RAPD variation (Volis *et al*, 2001, 2003) and quantitative traits (this paper). As a multiyear reciprocal introduction of seeds and seedlings conducted at one (pivot) location per region revealed local adaptation (Volis *et al*, 2002d), interpretation of the observed quantitative, allozyme and RAPD differentiation can be performed on the basis of a known effect of region-specific natural selection. We have the following objectives: (1) analysis of the extent and structure of quantitative trait variation; (2) a comparison of marker-specific (quantitative, allozyme and RAPD) estimates of regional genetic diversity, level and pattern of differentiation at two hierarchical levels; (3) interpretation of intermarker variation as due to selective or stochastic processes. Wild barley is a predominant selfing species that exhibits strong population differentiation in phenotypic traits (Snow and Brody, 1984), allozymes (Brown *et al*, 1978a), RAPDs (Baum *et al*, 1997), SSRs (Turpeinen *et al*, 2001) and AFLP markers (Pakniyat *et al*, 1997). High population subdivision due to selfing and genetic drift that is detectable with neutral markers can potentially mask population differentiation due to diversifying selection, making the Q<sub>ST</sub> *vs* F<sub>ST</sub> comparison inconclusive. Our major prediction was that overall estimates of

regional subdivision (a spatial scale where adaptation was found to occur) will be distinctly higher for quantitative than for either allozymes or RAPDs markers, while population differentiation within regions in  $Q_{ST}$  and  $F_{ST}$  will not differ.

## Materials and methods

### Study populations

*H. spontaneum* Koch is an annual, diploid grass with an estimated self-fertilization rate of 98% (Brown *et al.*, 1978b). In Israel, despite its mainly Mediterranean and Irano-Turanian distribution in steppe-like formations, wild barley penetrates into desert (<200 mm annual rainfall) and mountain (up to 1600 m elevation) environments where it has stable populations. The plant material used in this study consisted of 20 populations sampled in Israel in 1996, employing a nested sampling design. Each of five populations was collected in environments that were as similar as possible and comprised a group representative of one of the following environments/vegetation communities (in order of increasing mean annual rainfall and predictability): desert, semisteppe batha, Mediterranean grassland and mountain (D1-5, B1-5, G1-5 and M1-5, respectively – Figure 1). Relief,

slope exposition, vegetation and soil types were kept constant in sampling localities within a group (for a detailed description of the environments, see Volis *et al.*, 2002d). Populations comprising a group were  $\leq 5$  km (mountain) or  $\leq 20$  km (other three groups) from each other. All 20 populations were used in a study of population genetic structure, and one (pivot) population from each group was used in a comparative study of plant life histories and in a test of local adaptation (Volis *et al.*, 2002d). Environment-specific (region-specific) natural selection was explicit and detected *de facto* by reciprocal transplants of seeds and seedlings (Volis *et al.*, 2002d). Significant genotype-by-environment interactions and highest fitness of genotypes in indigenous environments (except batha genotypes that had highest fitness in the grassland transplant site) were found, indicating local adaptation. Clear evidence of local adaptation was also demonstrated for two ecotypes, desert and batha, in an earlier study (Volis *et al.*, 2002b).

### Quantitative traits


One mature spike per plant was collected in 1996 from approximately 50 plants in each of 20 populations. From all populations, the seeds from nine random chosen spikes (one seed per spike) were sown in a greenhouse. After 3 weeks, when plants were at the 2–3 leaf stage, they were transferred to 10 l buckets filled with a potting mixture of equal parts peat and vermiculite with a single plant/pot. Buckets were placed in a nethouse at the Institutes for Applied Research (Beer Sheva, Israel) that contained drip lines with 2 l/h drippers with one dripper placed in each bucket. A randomized block design was used with a block containing 20 plants (one plant per population). Owing to some mortality during the experiment, the number of plants per population ranged from six to nine plants, and number of plants per region from 36–45 plants.

In a second common garden experiment, we used the progeny of eight randomly chosen F1 plants per population. The number of populations was four, with one (pivot) population per region. Prior to the experiment, the mother plants were planted under uniform conditions in a greenhouse and four offspring per mother plant were used in the experiment. As wild barley is predominantly autogamous, the offspring of each mother plant can be considered as a genetically identical single genotype. Each of four offspring was randomly assigned to one of four blocks containing 32 plants.

The common garden experiments were conducted at the same time and the same set of quantitative traits was measured. After the appearance of reproductive tillers, the first three reproductive tillers of each plant were tagged with colored tags representing the first, second and third tiller in order of awn appearance and each tiller was measured for: number of spikelets per spike (NSP), tiller height (TH), flag and penultimate leaf length and width (FLL, PLL, FLW and PLW), spike length (SPL), awn length (AWL), number of nodes (NN), and the number of days to awning (DAW) were recorded. At senescence, mean spikelet weight (SWT) was obtained from the three tagged spikes.

### Allozymes and RAPDs

Nine enzyme systems encoding 15 loci were examined in 25 plants per population by starch gel electrophoresis



**Figure 1** Mean annual rainfall (mm), annual temperature and geographic distribution of sampling localities of wild barley in Israel: M1-5 = mountain; G1-5 = grassland; B1-5 = semisteppe batha; D1-5 = desert.

(for details see Volis *et al*, 2003). RAPD analysis was performed using DNA from 14 plants per population as described in Volis *et al* (2001).

### Quantitative genetic analyses

The structure of variation in phenotypic traits was analyzed after running nested ANOVA by partitioning the total variance among regions (df 3), among populations within a region (df 16) and among individuals within a population (df 120–180) (common garden 1) or among populations (df 3), among genotypes within a population (df 28) and within genotypes (df 96) (common garden 2). In common garden 1, variance was also partitioned into among- and within-population components for each region separately.

The data from common garden 1 (20 populations) allowed us to determine the among-population ( $\sigma_{\text{Pop}}^2$ ) and among-region ( $\sigma_{\text{Reg}}^2$ ) genetic variance, and to estimate regional subdivision  $Q_{\text{RT}}$  and population subdivision  $Q_{\text{ST}}$  as  $V_{\text{GB}}/(V_{\text{GB}} + V_{\text{GW}})$ , where  $V_{\text{GB}}$  is the proportion of genetic variance distributed either among populations ( $\sigma_{\text{Pop}}^2$ ) or among regions ( $\sigma_{\text{Reg}}^2$ ) and  $V_{\text{GW}}$  is the proportion of genetic variance distributed within populations (Spitze, 1993; Morgan *et al*, 2001). The population subdivision within each region ( $Q_{\text{SR}}$ ) was analyzed by conducting nested analysis of variance for each region separately. As genotypes were not replicated in the 20 population experiment, the within- and among-population components comprised both genetic and pure environmental components. However, we consider such  $Q_{\text{ST}}$  a useful surrogate for true  $Q_{\text{ST}}$  (with only a slight upward bias) because the environmental component for all traits except YLD was small and broad-sense heritability was very high (as revealed in the four population experiment; Table 2). For individual trait  $Q_{\text{RT}}$  values, the 95% confidence limits were calculated, adopting the equation given in Lynch and Walsh (1998) as

$$\frac{(F/F_U) - 1}{(F/F_U) + n - 1} < Q_{\text{RT}} < \frac{(F/F_L) - 1}{(F/F_L) + n - 1}$$

where  $F$ ,  $F_U$  and  $F_L$  are the observed, upper and lower  $F$  values, respectively, associated with  $(\alpha/2)$  at  $(N-1)$ ,  $(T-N)$  degrees of freedom. For the average  $Q_{\text{RT}}$  and  $Q_{\text{SR}}$ , the 95% confidence intervals were estimated by bootstrapping with random reassignment to regenerate the original sample sizes repeated 1000 times. This was done using the program RESAMPLING STATS (Simon, 1995).

Data obtained in common garden 2 (four populations) were used to estimate broad-sense heritability,  $h^2 = V_G/(V_G + V_E)$ . One-way ANOVA was performed for each of the four populations. The within-genotype variance ( $\sigma_E^2$ ) estimated the environmental variance  $V_E$  and the among-genotype variance provided an estimate of the genetic variance  $V_G$ . The standard errors of trait  $h^2$  were calculated following Lynch and Walsh (1998) as  $(1 - h^2)[1 + (n - 1)h^2]\sqrt{2/[Nn(n - 1)]}$ , where  $N$  is the number of genotypes and  $n$  is the number of individuals per genotype.

The amount of trait-specific regional variance was expressed as the coefficient of variation  $CV = \sqrt{\sigma_{\text{Reg}}^2}/X$ , where  $X$  is the regional phenotype mean, and compared

across regions by Levene's Test (JMP, 1995). The regional means were compared by using Welch ANOVA which allows variances to be unequal. The region differences over all traits considered simultaneously were analyzed by a single factor MANOVA followed by pairwise group contrasts. All the analyses of variance were done using JMP version 3.1.2 statistical package (JMP, 1995).

Principal component analysis was conducted on standardized original values using the Multi-Variate Statistical Package (MVSP, Kovach, 1999). A matrix of pairwise Euclidean distances between 20 populations was calculated by NTSYSpc program (Rohlf, 1998) and used in a Mantel test against RAPD and allozyme genetic distance matrices. Evaluation of significance of the matrix correlation employed a comparison of the observed Mantel test statistic,  $Z$ , with its random distribution generated by 1000 random permutations after the sequential Bonferroni adjustment for multiple testing.

### Molecular genetic analyses

For both RAPDs and allozymes, the distribution of genetic variability within and among populations and regions was investigated by an analysis of molecular variance (AMOVA; Excoffier *et al*, 1992) and using Nei's  $G_{\text{ST}}$  statistic (Nei, 1973). For the former, genetic distances between haplotypes were calculated based on squared Euclidean distances. As allozymes showed very low observed heterozygosity (<1%) (Volis *et al*, 2003), the heterozygotes were excluded and the data treated as haplotypes. For the latter, allele frequencies calculated from the RAPD bands assumed full dominance. Analyses were carried out for the whole sample and for each region separately. The number of permutations for significance testing was set at 1000 for analysis. The  $G_{\text{ST}}$  statistic was calculated with POPGENE version 1.31 (Yeh *et al*, 1998). This program was also used to calculate the pairwise population genetic distances (Nei, 1978) used in a Mantel test against each other and genetic distances calculated from quantitative traits. Principal component analysis was used to ordinate relationships among individuals and among regions using the MVSP program.

## Results


### Quantitative genetic variation

Regional means and coefficients of variation for 11 quantitative traits are shown in Table 1. Regional values differed significantly in each trait (Welch ANOVA) and over all traits analyzed simultaneously (Wilks'  $\lambda = 0.04$ ,  $F_{33,430} = 26.8$ ,  $P < 0.001$ , MANOVA). MANOVA also revealed highly significant differences ( $P < 0.001$ ) in each pairwise region comparison. Variation within groups, although different for some traits (as revealed by Levene's test), showed no consistent pattern across the groups. In all traits, we found a trend of either an increase or decrease in trait values from desert to grassland, with plants from the mountain region strongly resembling those from the grassland (Table 1 and Figure 2). Heritability of all traits except YLD was high, ranging 0.33–0.94 when averaged over four populations, but with striking differences among populations

**Table 1** Regional means and coefficients of variation (in parentheses) for 11 quantitative traits

Region	n	DAW	NSP	SWT	YLD	SPL	AWL	FLL	FLW	PLL	PLW	TH
Desert	36	120.0 (3.8)	37.4 (8.6)	48.6 (14.6)	2884 (26.3)	20.3 (7.9)	6.0 (16.7)	13.4 (24.6)	1.09 (15.6)	26.8 (18.7)	1.51 (12.6)	63.9 (13.1)
Batha	44	131.4 (3.9)	33.9 (8.8)	70.3 (18.9)	2444 (20.8)	24.3 (7.0)	10.1 (16.8)	9.2 (27.2)	0.90 (20.0)	17.8 (20.2)	1.42 (21.8)	77.6 (11.2)
Grassland	45	140.3 (3.6)	29.1 (7.9)	79.4 (13.6)	2438 (18.8)	28.6 (11.9)	15.2 (22.4)	6.0 (23.3)	0.53 (22.6)	13.2 (17.4)	1.00 (15.0)	85.7 (13.0)
Mountain	39	145.3 (2.8)	31.6 (9.5)	70.4 (11.2)	2498 (16.8)	28.9 (11.1)	14.0 (20.0)	7.2 (26.4)	0.51 (21.6)	14.8 (16.9)	0.89 (18.0)	87.4 (9.6)
Levene's test		0.94 ns	1.95 ns	3.8*	4.0**	12.5***	15.6***	8.0***	2.9*	10.3***	8.3***	1.7 ns
Welch's ANOVA		234.1***	62.8***	96.0***	3.4*	118.0***	179.0***	62.2***	57.8***	81.8***	96.7***	56.4***

Level of significance: \* – 0.05; \*\* – 0.01; \*\*\* – 0.001; ns = not significant.



**Figure 2** Plot of principal components for quantitative trait variation.

(Table 2). Average trait  $h^2$  was significantly higher in the desert population than in batha and mountain populations (Wilcoxon Signed-Ranks Test,  $Z = 2.49$  and  $1.96$ ,  $P < 0.05$ ), and higher in the grassland population than in batha (Wilcoxon Signed-Ranks Test,  $Z = 2.40$ ,  $P < 0.05$ ). The traits can be divided into two groups on the basis of their intrapopulation differences in heritability: (1) with uniformly high ( $> 0.8$ ) or low ( $< 0.3$ )  $h^2$  (SPL, AWL and YLD); (2) with highly variable  $h^2$  among populations (remaining traits).

Partitioning of variation in quantitative traits revealed a very high regional component and regional subdivision  $Q_{RT}$  ranged from 0.55 to 0.89 for all traits except one (YLD) (Table 2). Regions contributed more to the total variation than populations, genotypes or environment altogether in all traits except YLD and TH. Subdivision among populations within regions averaged over traits ranged from 0.17 to 0.40 (Table 3).

#### Molecular variation for RAPDs and allozymes

Total and within-population gene diversity estimates ( $H_T$  and  $H_S$ ) for RAPDs and allozymes averaged across loci did not differ, nor did four regional estimates (Table 3). Despite the fact that the patterns of differentiation in RAPDs and allozymes matched in general, the allozymes exhibited lower regional subdivision than RAPDs (Table 3), and most of this variation appears to be due to segregation of the mountain population from the rest (not shown), with variation in allozymes being much more discontinuous than in RAPDs. The latter appears to be due to a limited number of polymorphic allozyme loci (only five loci exhibited  $H_T > 0.3$ ). The mountain region was clearly distinct for both marker types but segregation of the other three regions was less obvious, especially for allozymes (not shown). The levels of regional subdivision and population subdivision within regions were similar for both types of markers with high agreement between the estimates of subdivision by AMOVA and Nei's gene diversity (Table 3).

#### Relationships between molecular and quantitative trait variation

A test of the relationships between three matrices of pairwise genetic distances (quantitative trait values,


**Table 2** Estimates of variance components (in %), regional differentiation (Q<sub>RT</sub>) and heritability (h<sup>2</sup>) for 11 quantitative traits

Trait	20 populations experiment										4 populations experiment									
	Variance components					Q <sub>RT</sub>					Variance components					h <sup>2</sup>				
	σ <sup>2</sup> <sub>Reg</sub>	σ <sup>2</sup> <sub>Pop</sub>	σ <sup>2</sup> <sub>Within</sub>	σ <sup>2</sup> <sub>Gen</sub>		σ <sup>2</sup> <sub>E</sub>		σ <sup>2</sup> <sub>Reg+Pop</sub>		σ <sup>2</sup> <sub>E</sub>		Desert		Batha		Grassland		Mountain		
DAW	81.4	9.3	9.3	0.897 (0.857–0.965)	62.9	34.4	2.7	0.972 (0.008)	0.689 (0.069)	0.830 (0.043)	0.797 (0.050)									
NSP	53.8	17.1	29.1	0.649 (0.531–0.856)	32.4	52.7	14.9	0.725 (0.063)	0.492 (0.091)	0.795 (0.050)	0.912 (0.024)									
SWT	56.1	10.8	33.1	0.629 (0.492–0.838)	70.8	8.7	20.5	0.778 (0.053)	0.118 (0.086)	0.427 (0.094)	-0.017 (0.070)									
YLD	4.8	18.7	76.5	0.059 (-0.186–0.142)	28.4	21.2	64.5	0.022 (0.075)	0.077 (0.082)	0.242 (0.094)	0.105 (0.085)									
SPL	65.5	17.4	17.1	0.793 (0.722–0.926)	77.2	18.7	1.6	0.973 (0.008)	0.938 (0.017)	0.817 (0.045)	0.921 (0.022)									
AWL	68.5	17.0	14.5	0.825 (0.762–0.938)	80.2	43.7	1.0	0.985 (0.004)	0.961 (0.011)	0.924 (0.020)	0.900 (0.027)									
FLI	60.6	6.0	33.4	0.645 (0.514–0.848)	43.0	20.0	13.2	0.813 (0.046)	0.403 (0.095)	0.807 (0.048)	0.262 (0.095)									
FLW	76.6	5.9	17.5	0.814 (0.738–0.931)	64.4	37.4	15.5	0.698 (0.067)	0.211 (0.093)	0.603 (0.080)	-0.011 (0.071)									
PLL	72.1	2.7	25.2	0.741 (0.640–0.898)	35.0	18.7	27.5	0.802 (0.049)	0.224 (0.094)	0.176 (0.091)	0.523 (0.088)									
PLW	61.8	17.2	21	0.746 (0.658–0.904)	65.2	11.0	16.1	0.512 (0.089)	0.517 (0.089)	0.684 (0.070)	0.118 (0.086)									
TH	47.8	13.2	39	0.551 (0.401–0.794)	80.9	7.0	8.0	0.661 (0.073)	0.156 (0.089)	0.574 (0.084)	0.737 (0.061)									
Mean CI 95% (bootstrapped over traits)	59.0	12.3	28.7	0.668 (0.471–0.757)	58.2	24.9	16.9	0.722 (0.495–0.840)	0.435 (0.277–0.645)	0.626 (0.462–0.755)	0.477 (0.249–0.691)									

Variance components: σ<sup>2</sup><sub>Reg</sub> = among-region; σ<sup>2</sup><sub>Pop</sub> = among-population; σ<sup>2</sup><sub>Gen</sub> = among-genotype; σ<sup>2</sup><sub>E</sub> = environmental; σ<sup>2</sup><sub>Reg+Pop</sub> = among-region including among-population component.

**Table 3** Molecular and quantitative trait subdivision estimated with θ-statistics (AMOVA; Excoffier et al, 1992), G<sub>ST</sub>-statistics (Spitze, 1993). Estimates of θ<sub>ST</sub> and G<sub>ST</sub> (RAPDs and allozymes) and Q<sub>ST</sub> (quantitative traits) are presented as both regional subdivision (θ<sub>RT</sub>, G<sub>RT</sub> and Q<sub>RT</sub>) and population subdivision within regions (θ<sub>SR</sub>, G<sub>SR</sub> and Q<sub>SR</sub>). Estimates of H<sub>T</sub>, Q<sub>RT</sub> and Q<sub>SR</sub> are provided with 95% CI (bootstrapping over loci/traits)

	RAPDs					Allozymes					Quantitative				
	θ <sub>KT</sub>	θ <sub>SR</sub>	H <sub>T</sub>	G <sub>RT</sub>	G <sub>SR</sub>	θ <sub>RT</sub>	θ <sub>SR</sub>	H <sub>T</sub>	G <sub>RT</sub>	G <sub>SR</sub>	Q <sub>RT</sub>	Q <sub>SR</sub>	Q <sub>RT</sub>	Q <sub>SR</sub>	
Total	0.205	0.256	0.253 (0.223–0.285)	0.207	0.311	0.162	0.096	0.224 (0.154–0.294)	0.108	0.112	0.628 (0.471–0.757)	0.324 (0.239–0.417)			
Desert		0.219	0.200 (0.169–0.247)		0.231		0.041	0.217 (0.163–0.276)		0.064		0.169 (0.114–0.256)			
Batha		0.166	0.207 (0.169–0.239)		0.175		0.092	0.193 (0.138–0.255)		0.086		0.261 (0.152–0.385)			
Grassland		0.378	0.202 (0.162–0.238)		0.355		0.144	0.207 (0.129–0.286)		0.168		0.274 (0.136–0.404)			
Mountain		0.509	0.194 (0.156–0.244)		0.494		0.177	0.184 (0.114–0.262)		0.134		0.396 (0.254–0.559)			



**Figure 3** Histograms of regional differentiation estimates for 54 RAPD loci ( $G_{RT}$ ), 17 allozyme loci ( $G_{RT}$ ) and 11 quantitative traits ( $Q_{RT}$ ) (from top to bottom).

allozyme and RAPD allele frequencies) calculated from 20 populations revealed the following: There was a highly significant relationship between allozyme and RAPD Nei's genetic distances ( $r=0.69$ ,  $P<0.001$ , Mantel test), but not between quantitative traits and either allozymes and RAPDs ( $r=0.02$  and  $0.26$ ,  $P>0.05$ , Mantel's test). The pairwise estimates of population subdivision  $\theta_{ST}$  for two molecular marker types were also highly correlated ( $r=0.70$ ,  $P<0.001$ , Mantel test) with one another. The within-population levels of genetic variation for quantitative traits (CV), allozymes and RAPDs  $H_E$  were not correlated with one another. The within-population gene diversity estimated with allozymes (mean = 0.173, range 0.114–0.253) was higher than that estimated with RAPDs (mean = 0.138, range 0.040–0.202) (Wilcoxon Signed-Ranks Test,  $Z=2.65$ ,  $P=0.008$ ).

The frequency distribution of regional subdivision estimates was skewed in opposite directions for quantitative traits (mode  $0.6 < Q_{RT} < 0.9$ ) vs allozyme and RAPD loci (mode  $0 < G_{RT} < 0.1$ ) (Figure 3). The allozyme loci  $G_{RT}$  values, however, were almost exclusively located near the mode, while RAPD  $G_{RT}$  estimates varied widely.

## Discussion

### Quantitative genetic variation

Levels of regional subdivision for quantitative traits ( $Q_{RT}$ ) were very high (0.55–0.90 in 10 out of 11 traits). These traits were previously analyzed for phenotypic

plasticity (Volis *et al.*, 2002e) and involvement in the adaptive process (Volis *et al.*, 2004), including involvement in region-specific adaptation (Volis *et al.*, 2002c), using plants from one (pivot) population per region. In these experiments, DAW, SPL, SWT, AWL and TH were significant predictors of individual plant fitness. Leaf dimensional traits (FFL, FLW, PLL, PLW) have also shown clear regional differences, apparently indicating region-specific adaptation (Volis *et al.*, 2002e). The results reported on here were obtained using plants from 20 populations and are in full agreement with those previously derived from pivot populations only. Together, they suggest a region-specific adaptive process in wild barley that is detectable with quantitative traits. There is directional selection in *H. spontaneum* along a gradient of increasing aridity (from mountain to desert) for early transition to reproduction, and along the same aridity gradient from grassland to desert for increase in size of flag and penultimate leaves and number of seeds per spike, and a decrease in seed weight and size of reproductive structures. The plants from mountain experience specific environmental conditions (longer vegetative period and much colder winter with occasional frosts) but adaptation to these conditions may be achieved by physiological and not morphological traits, as the latter were very similar to those of grassland plants.

Regional genetic differentiation, both clinal and non-clinal, observed in studied quantitative traits is clearly adaptive after several reciprocal introductions performed either at two (Volis *et al.*, 2002b) or four (Volis *et al.*, 2002d) sampling locations. However, Verhoeven *et al.* (2004), who demonstrated local adaptation to different nutrient environments in wild barley, failed to detect diversifying selection on the underlying QTL loci. The results of Verhoeven *et al.* (2004) show that the common view that adaptive differentiation involves a genetic trade-off between alleles of the same locus that have opposite fitness effects in contrasting environments needs to be reconciled with an alternative view that adaptive differentiation is caused by loci that differ in the magnitude but not in the direction of their fitness effects across habitats (Kawecki, 1997).

### Molecular variation for RAPDs and allozymes and relationship between markers

It has been predicted that the extent of genetic variation in quantitative traits and molecular gene diversity (expressed as expected heterozygosity  $H_E$ ) are correlated (Falconer and Mackay 1996). This, however, should be true only if the variation is neutral, because such forces as drift and migration will similarly affect neutral but not fitness-related loci (Lewontin and Krakauer, 1973; Slatkin, 1987). As the quantitative traits examined in this study were found to be adaptive, it is not surprising that we failed to detect a relationship between either allozyme or RAPD  $H_E$  and variation in any of the quantitative traits. A lack of correlation between within-population allozyme and quantitative variation was also found in two other studies performed on wild plant species (Waldmann and Andersson, 1998; Podolsky, 2001).

Our results indicate that there is a positive relationship between interpopulation genetic distances obtained from

allozyme- and RAPD-based data, which is consistent with the results of many other studies (eg Heun *et al*, 1994; Lifante and Aguinalgalde, 1996; Wu *et al*, 1999; Bartish *et al*, 2000a). However, we did not find a positive relationship between corresponding estimates of within-population variation. In contrast to generally observed similarities in allozyme- and RAPD-generated estimates of within-population variation or higher estimates for RAPDs (reviewed in Vandewoestijne and Baguette, 2002; Nybom, 2004), we found the opposite pattern. Higher regional and population subdivision for RAPDs *vs* allozymes contrasted with lower within-population variation for RAPDs. One possible explanation could be a smaller sample size for RAPDs *vs* allozymes (14 and 25 plants per population, respectively).

Patterns of inter-regional variation for quantitative traits and the two molecular markers visualized by principal components analysis were not congruent. However, allozyme and RAPD scatterplots showed more in common than did a scatterplot of quantitative traits with either of the other two (in particular, both molecular marker types, but not quantitative traits, revealed a segregation of the mountain region from the other three regions). Additionally, matrices of allozyme and RAPD pairwise population genetic distances were significantly correlated with each other but not with a matrix for quantitative traits. As quantitative traits were found to be adaptive, this implies that most allozyme and RAPD variation is nonadaptive.

The degree of regional subdivision for allozymes and RAPDs was lower than for quantitative traits. The level of regional subdivision in two molecular markers was comparable with population subdivision within a region, while in quantitative traits it was significantly (almost 2 ×) higher. Disagreement between the structure of variation at two scales (regions and populations within regions) in quantitative traits and two molecular markers, and the pattern of molecular variation reported here and in Volis *et al* (2001, 2003) point at selective neutrality of overall allozyme and RAPD variation. However, effects of selection or gene hitch-hiking on allozyme and RAPD variation cannot be ruled out completely. The frequency distributions of G<sub>RT</sub> values of allozyme and RAPD loci show that most loci exhibit minor or no regional differentiation, and that the relatively high overall estimates of G<sub>RT</sub> are due to several loci with exceptionally high regional differentiation. This pattern is difficult to explain by genetic drift alone as all neutral loci are expected to be affected in the same manner. Natural selection, operating on these exceptional loci, is a more plausible scenario. However, although this scenario is plausible, it still needs to be supported by experiments. An alternative explanation of nonuniformity of genetic subdivision across loci is that the selective process has an effect on those neutral loci that are in linkage disequilibrium with loci directly affected by selection (Maynard Smith and Haigh, 1974). This stems from the theory of hitch-hiking that, within a single population and under homogeneous environmental conditions, reduced gene diversity is expected at a neutral locus closely linked to the selected locus (Maynard Smith and Haigh, 1974; Barton, 1995). However, in a geographically subdivided population (Slatkin and Wiehe, 1998) or under local adaptation (Charlesworth *et al*, 1997), hitch-hiking may lead to high G<sub>ST</sub>

values even in the case of significant gene flow, and despite simultaneous reduction in within-population gene diversity.

In conclusion, the results of this study showed a moderate to high degree of differentiation among populations of wild barley in both molecular (allozymes and RAPDs) and quantitative traits. However, examination of population subdivision at two spatial scales, one of which was ecologically relevant, revealed two different outcomes. The Q<sub>ST</sub> and F<sub>ST</sub> did not differ at the population scale, that is, within regions, where stochastic evolutionary forces appear to prevail, but differed at the regional scale, where an adaptive differentiation process was previously found to occur. This implies that the comparison of levels of subdivision in quantitative traits (most of which are closely related to fitness) with molecular markers (which fitness effects are unknown) is indicative of adaptive population differentiation only when sampling is done at the proper scale.

## Acknowledgements

We thank Richard Nichols and two anonymous reviewers for their helpful comments. A grant from the Israel Academy of Sciences (86293101) supported this study.

## References

- Aagaard J, Krutovskii KV, Strauss SH (1998). RAPDs and allozymes exhibit similar levels of diversity and differentiation among populations and races of Douglas-fir. *Heredity* **81**: 69–78.
- Allendorf FW, Seeb LW (2000). Concordance of genetic divergence among sockeye salmon populations at allozyme, nuclear DNA, and mitochondrial DNA markers. *Evolution* **54**: 640–651.
- Ayres DR, Ryan FJ (1997). The clonal and population structure of a rare endemic plant, *Wyethia reticulata* (Asteraceae): allozyme and RAPD analysis. *Mol Ecol* **6**: 761–772.
- Bartish IV, Garkava LP, Rumpunen K, Nybom H (2000a). Phylogenetic relationships and differentiation among and within populations of *Chaenomeles* Lindl. (Rosaceae) estimated with RAPDs and isozymes. *Theoret Appl Genet* **101**: 554–563.
- Bartish IV, Jeppsson N, Bartish GI, Lu R, Nybom H (2000b). Inter- and intraspecific genetic variation in *Hippophae* (Elaeagnaceae) investigated by RAPD markers. *Plant Syst Evol* **225**: 85–101.
- Barton NH (1995). Linkage and the limits to natural selection. *Genetics* **140**: 821–841.
- Baum BR, Nevo E, Johnson DA, Beiles A (1997). Genetic diversity in wild barley (*Hordeum spontaneum* C. Koch) in the Near East: a molecular analysis using Random Amplified Polymorphic DNA (RAPD) markers. *Genet Resources Crop Evol* **44**: 147–157.
- Black-Samuelsson S, Eriksson G, Gustafsson L, Gustafsson P (1997). RAPD and morphological analysis of the rare plant species *Vicia pisiformis* (Fabaceae). *Biol J Linnean Soc* **61**: 325–343.
- Bonnin I, Prospero JM, Olivieri I (1996). Genetic markers and quantitative genetic variation in *Medicago truncatula* (Leguminosae): a comparative analysis of population structure. *Genetics* **143**: 1795–1805.
- Bossart JL, Pashley Prowell D (1998). Genetic estimates of population structure and gene flow: limitations, lessons and new directions. *Trends Ecol Evol* **13**: 202–206.

- Brown AHD, Nevo E, Zohary D, Dagan O (1978a). Genetic variation in natural populations of wild barley (*Hordeum spontaneum*). *Genetica* **49**: 97–108.
- Brown AHD, Zohary D, Nevo E (1978b). Outcrossing rates and heterozygosity in natural populations of *Hordeum spontaneum* Koch in Israel. *Heredity* **41**: 49–62.
- Cavalli-Sforza L (1966). Population structure and human evolution. *Proc R Soc London, Ser B* **164**: 362–379.
- Chan Y, Arcese P (2003). Morphological and microsatellite differentiation in *Melospiza melodia* (Aves) at a microgeographic scale. *J Evol Biol* **16**: 939–947.
- Charlesworth D (2003). Effects of inbreeding on the genetic diversity of populations. *Philos Trans R Soc London Ser B-Biol Sci* **358**: 1051–1070.
- Charlesworth B, Nordborg M, Charlesworth D (1997). The effects of local selection, balanced polymorphism and background selection on equilibrium patterns of genetic diversity in subdivided populations. *Genet Res* **70**: 155–174.
- Comes HP, Abbott RJ (2000). Random amplified polymorphic DNA (RAPD) and quantitative trait analyses across a major phylogeographic break in the Mediterranean ragwort *Senecio gallicus* (Asteraceae). *Mol Ecol* **9**: 61–76.
- Edmunds S, Harrison JS (2003). Molecular and quantitative trait variation within and among populations of the intertidal copepod *Tigriopus californicus*. *Evolution* **57**: 2277–2285.
- Excoffier L, Smouse PE, Quattro JM (1992). Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* **131**: 479–491.
- Falconer DS, Mackay TFC (1996). *Introduction to Quantitative Genetics*, 4th edn. Longman: Harlow, UK.
- Heaton HJ, Whitkus R, Gomez PA (1999). Extreme ecological and phenotypic differences in the tropical tree chicozapote (*Manilkara zapota* (L.) P. Royen) are not matched by genetic divergence: a random amplified polymorphic DNA (RAPD) analysis. *Mol Ecol* **8**: 627–632.
- Heun M, Murphy JP, Phillips TD (1994). A comparison of RAPD and isozyme analyses for determining the genetic relationships among *Avena sterilis* L. accessions. *Theoret Appl Genet* **87**: 689–696.
- Isabel N, Beaulieu J, Bousquet J (1995). Complete congruence between gene diversity estimates derived from genotypic data at enzyme and random amplified polymorphic DNA loci in black spruce. *PNAS* **92**: 6369–6373.
- Jaramillo-Correa JP, Beaulieu J, Bousquet J (2001). Contrasting evolutionary forces driving population structure at expressed sequence tag polymorphisms, allozymes and quantitative traits in white spruce. *Mol Ecol* **10**: 2729–2740.
- Jenczewski E, Prospero JM, Ronfort J (1999). Differentiation between natural and cultivated populations of *Medicago sativa* (Leguminosae) from Spain: Analysis with random amplified polymorphic DNA (RAPD) markers and comparison to allozymes. *Mol Ecol* **8**: 1317–1330.
- JMP (1995). SAS Institute Inc. Cary, NC.
- Karhu A, Hurme P, Karjalainen M, Karvonen P, Kärkkäinen K, Neale D *et al* (1996). Do molecular markers reflect pattern of differentiation in adaptive traits of conifers? *Theoret Appl Genet* **93**: 215–221.
- Karl SA, Avise JC (1992). Balancing selection at allozyme loci in oysters: implications from nuclear RFLPs. *Science* **256**: 100–102.
- Kawecki TJ (1997). Sympatric speciation via habitat specialization driven by deleterious mutations. *Evolution* **51**: 1751–1763.
- Kimura M (1983). The neutral theory of molecular evolution. In: Mei M, Koehn RK (eds) *Evolution of Genes and Proteins*. Sinauer Associates: Sunderland, MA. pp 208–233.
- Knapp EE, Rice KJ (1998). Comparison of isozymes and quantitative traits for evaluating patterns of genetic variation in purple needlegrass (*Nassella pulchra*). *Conservation Biol* **12**: 1031–1041.
- Kovach W (1999). *Multi-Variate Statistical Package 3.11b*. Kovach Computing Services: Aberystwyth, Wales, UK.
- Kuittinen H, Mattila A, Savolainen O (1997). Genetic variation at marker loci and in quantitative traits in natural populations of *Arabidopsis thaliana*. *Heredity* **79**: 144–152.
- Lagercrantz U, Ryman N (1990). Genetic structure of Norway spruce (*Picea abies*): concordance of morphological and allozymic variation. *Evolution* **44**: 38–53.
- Latta RG, Mitton JB (1997). A comparison of population differentiation across four classes of gene marker in limber pine (*Pinus flexilis* James). *Genetics* **146**: 1153–1163.
- Le Corre V, Dumolin-Lapegue S, Kremer A (1997). Genetic variation at allozyme and RAPD loci in sessile oak *Quercus petraea* (Matt) Liebl: the role of history and geography. *Mol Ecol* **6**: 519–529.
- Lewontin RC (1991). Electrophoresis in the development of evolutionary genetics: milestone or millstone? *Genetics* **128**: 657–662.
- Lewontin RC, Krakauer J (1973). Distribution of gene frequency as a test of the theory of the selective neutrality of polymorphisms. *Genetics* **74**: 175–195.
- Lifante ZD, Aguinalgalde I (1996). The use of random amplified polymorphic DNA (RAPD) markers for the study of taxonomical relationships among species of *Asphodelus* sect. *Verinea* (Asphodelaceae). *Am J Botany* **83**: 949–953.
- Luttikhuisen PC, Drent J, van Delden W, Piersma T (2003). Spatially structured genetic variation in a broadcast spawning bivalve: quantitative vs molecular traits. *J Evol Biol* **16**: 260–272.
- Lynch M (1988). The divergence of neutral quantitative characters among partially isolated populations. *Evolution* **42**: 455–466.
- Lynch M (1994). The neutral theory of phenotypic evolution. In: Real L (ed) *Ecological Genetics*. Princeton University Press: Princeton, NJ. pp 86–108.
- Lynch M, Walsh B (1998). *Genetics and Analysis of Quantitative Traits*. Sinauer Associates: Sunderland, MA.
- Mamuris Z, Stamatis C, Triantaphyllidis C (1999). Intraspecific genetic variation of striped red mullet (*Mullus surmuletus* L.) in the Mediterranean Sea assessed by allozyme and random amplified polymorphic DNA (RAPD) analysis. *Heredity* **83**: 30–38.
- Maynard Smith J, Haigh J (1974). The hitch-hiking effect of a favorable gene. *Genet Res* **23**: 23–35.
- McDonald JH (1994). Detecting natural selection by comparing geographic variation in protein and DNA polymorphisms. In: Golding B (ed) *Non-Neutral Evolution. Theories and Molecular Data*. Chapman & Hall: New York. pp 88–100.
- McKay JK, Bishop JG, Lin JZ, Richards JH, Sala A, Mitchell-Olds T (2001). Local adaptation across a climatic gradient despite small effective population size in the rare sapphire rockcress. *Proc R Soc London Ser B-Biol Sci* **268**: 1715–1721.
- Merila J, Crnokrak P (2001). Comparison of genetic differentiation at marker loci and quantitative traits. *J Evol Biol* **14**: 892–903.
- Morgan KK, Hicks J, Spitze K, Latta L, Pfrender ME, Weaver CS *et al* (2001). Patterns of genetic architecture for life-history traits and molecular markers in a subdivided species. *Evolution* **55**: 1753–1761.
- Nei M (1973). Analysis of gene diversity in subdivided populations. *Proc Natl Acad Sci* **70**: 3321–3323.
- Nei M (1978). Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics* **89**: 583–590.
- Nybom H (2004). Comparison of different nuclear DNA markers for estimating intraspecific genetic diversity in plants. *Mol Ecol* **13**: 1143–1155.
- Pakniyat H, Powell W, Baird E, Handley LL, Robinson D, Scrimgeour CM *et al* (1997). AFLP variation in wild barley (*Hordeum spontaneum*). *Genome* **40**: 332–341.

- Peakall R, Smouse PE, Huff DR (1995). Evolutionary implications of allozyme and RAPD variation in diploid populations of dioecious buffalograss *Buchloe dactyloides*. *Mol Ecol* **4**: 135–147.
- Petit C, Freville H, Mignot A, Colas B, Riba M, Imbert E *et al* (2001). Gene flow and local adaptation in two endemic plant species. *Biol Conservation* **100**: 21–34.
- Podolsky RH (2001). Genetic variation for morphological and allozyme variation in relation to population size in *Clarkia dudleyana*, an endemic annual. *Conservation Biol* **15**: 412–423.
- Podolsky RH, Holtsford TP (1995). Population structure of morphological traits in *Clarkia dudleyana*. 1. Comparison of F<sub>st</sub> between allozymes and morphological traits. *Genetics* **140**: 733–744.
- Rogers AR, Harpending HC (1983). Population structure and quantitative characters. *Genetics* **105**: 985–1002.
- Rohlf FJ (1998). *NTSYSpc Numerical Taxonomy and Multivariate Analysis System, version 2.0*. Eeter Software: Setauket, New York.
- Ross KG (1997). Multilocus evolution in fire ants: effects of selection, gene flow and recombination. *Genetics* **145**: 961–974.
- Ross KG, Krieger MJB, Shoemaker DD, Vargo EL, Keller L (1997). Hierarchical analysis of genetic structure in native fire ant populations: results from three classes of molecular markers. *Genetics* **147**: 643–655.
- Ross KG, Shoemaker DD, Krieger MJB, DeHeer CJ, Keller L (1999). Assessing genetic structure with multiple classes of molecular markers: a case study involving the introduced fire ant *Solenopsis invicta*. *Mol Biol Evol* **16**: 525–543.
- Simon JL (1995). Resampling: the New Statistics. Julian L Simon.
- Slatkin M (1987). Gene flow and the geographic structure of natural populations. *Science* **236**: 787–792.
- Slatkin M, Wiehe T (1998). Genetic hitch-hiking in a subdivided population. *Genet Res* **71**: 155–160.
- Snow L, Brody T (1984). Genetic variation of *Hordeum spontaneum* in Israel: eco-geographical races, detected by trait measurements. *Plant Syst Evol* **145**: 15–28.
- Sokal RR, Wartenburg DE (1983). A test of spatial autocorrelation analysis using an isolation-by-distance model. *Genetics* **105**: 219–237.
- Spitze K (1993). Population structure in *Daphnia obtusa*: quantitative genetic and allozymic variation. *Genetics* **135**: 367–374.
- Steinger T, Haldimann P, Leiss KA, Muller-Scharer H (2002). Does natural selection promote population divergence? A comparative analysis of population structure using amplified fragment length polymorphism markers and quantitative traits. *Mol Ecol* **11**: 2583–2590.
- Storz JF (2002). Contrasting patterns of divergence in quantitative traits and neutral DNA markers: analysis of clinal variation. *Mol Ecol* **11**: 2537–2551.
- Sun M, Wong KC (2001). Genetic structure of three orchid species with contrasting breeding systems using RAPD and allozyme markers. *Am J Botany* **88**: 2180–2188.
- Szmidt AE, Wang XR, Lu MZ (1996). Empirical assessment of allozyme and RAPD variation in *Pinus sylvestris* (L) using haploid tissue analysis. *Heredity* **76**: 412–420.
- Turpeinen T, Tenhola T, Manninen O, Nevo E, Nissila E (2001). Microsatellite diversity associated with ecological factors in *Hordeum spontaneum* populations in Israel. *Mol Ecol* **10**: 1577–1591.
- Vandewoestijne S, Baguette M (2002). The genetic structure of endangered populations in the Cranberry Fritillary, *Boloria aquilonaris* (Lepidoptera, Nymphalidae): RAPDs vs allozymes. *Heredity* **89**: 439–445.
- Verhoeven KJF, Vanhala TK, Biere A, Nevo E, Van Damme JMM (2004). The genetic basis of adaptive population differentiation: a quantitative trait locus analysis of fitness traits in two wild barley populations from contrasting habitats. *Evolution* **58**: 270–283.
- Volis S, Mendlinger S, Turuspekov Y, Esnazarov U (2002a). Phenotypic and allozyme variation in Mediterranean and desert populations of wild barley, *Hordeum spontaneum* Koch. *Evolution* **56**: 1403–1415.
- Volis S, Mendlinger S, Ward D (2002b). Adaptive traits of wild barley plants of Mediterranean and desert origin. *Oecologia* **133**: 131–138.
- Volis S, Mendlinger S, Ward D (2002c). Differentiation along a gradient of environmental productivity and predictability in populations of *Hordeum spontaneum* Koch: multilevel selection analysis. *Biol J Linnean Soc* **75**: 313–318.
- Volis S, Mendlinger S, Ward D (2002d). Differentiation in populations of *Hordeum spontaneum* along a gradient of environmental productivity and predictability: life history and local adaptation. *Biol J Linnean Soc* **77**: 479–490.
- Volis S, Mendlinger S, Ward D (2002e). Differentiation in populations of *Hordeum spontaneum* Koch along a gradient of environmental productivity and predictability: plasticity in response to water and nutrient stress. *Biol J Linnean Soc* **75**: 301–312.
- Volis S, Shulgina I, Ward D, Mendlinger S (2003). Regional allozyme variation in wild barley, *Hordeum spontaneum*: adaptive or neutral? *J Hered* **94**: 341–351.
- Volis S, Verhoeven K, Mendlinger S, Ward D (2004). Phenotypic selection and regulation of reproduction in different environments in wild barley. *J Evol Biol* **17**: 1121–1131.
- Volis S, Yakubov B, Shulgina I, Ward D, Zur V, Mendlinger S (2001). Tests for adaptive RAPD variation in population genetic structure of wild barley, *Hordeum spontaneum* Koch. *Biol J Linnean Soc* **74**: 289–303.
- Waldmann P, Andersson S (1998). Comparison of quantitative genetic variation and allozyme diversity within and between populations of *Scabiosa canescens* and *S. columbaria*. *Heredity* **81**: 79–86.
- Wu J, Krutovskii KV, Strauss SH (1999). Nuclear DNA diversity, population differentiation, and phylogenetic relationships in the California closed-cone pines based on RAPD and allozyme markers. *Genome* **42**: 893–908.
- Yang RC, Yeh FC, Yanchuk A (1996). A comparison of isozyme and quantitative genetic variation in *Pinus contorta* ssp. *latifolia* by F<sub>ST</sub>. *Genetics* **142**: 1045–1052.
- Yeh FC, Rong-Cai Y, Boyle T (1998). *POPGENE VERSION 1.31*. University of Alberta, Center for International Forestry Research: Edmonton, Alberta, Canada.