

Inflationary cosmology in the central region of string/M-theory moduli Space

Ram Brustein




אוניברסיטת בן-גוריון

hep-th/0205042
hep-th/02xxxx

with S. de Alwis and E. Novak
PRL 87 (2001), hep-th/0106174
PRD 64 (2001), hep-th/0002087
with S. de Alwis

- * Outer region of moduli space: problems!
- * Central region: parametrization with $N=1$ SUGRA
- * Scales & shape of central region potential
- * Inflation: constraints & predictions

String Moduli Space



Requirements

- $D=4$
- $N=1$ SUSY $\rightarrow N=0$
- $CC < (m_{3/2})^4$
- SM (will not discuss)
- Volume/Coupling moduli

T S

Perturbative theories =
cosmological disaster

- massless moduli
- Gravity \neq Einstein’s
- Inflation blocked

Central Region

Our proposal:

- Parametrization with $D=4$, $N=1$ SUGRA
- Stabilization by SSB effects @ string scale
- Continuously adjustable parameter
- SUSY breaking @ lower scale by FT effects
- PCCP o.k after SUSY breaking



Scales & Shape of Moduli Potential

- The width of the central region

In effective 4D theory:

moduli kinetic terms multiplied by $M_{\text{S}}^8 V_6$ ($M_{11}^9 V_7$ in M-theory).

Curvature term multiplied by same factors

“Calibrate” using 4D Newton’s constant $8\pi G_{\text{N}}=m_{\text{p}}^{-2}$

$$\Gamma = \frac{1}{2} \int d^4 x \left\{ \underline{m_p^2} \sqrt{-g} R + \underline{m_p^2} \partial_\mu \psi \partial^\mu \psi \right\}$$

→ Typical distances are order one in units of m_{p}

- The scale of the potential

$$W \approx M_S^3 \Rightarrow V \approx M_S^6 / m_p^2 \equiv \Lambda^4 \quad \text{NO VOLUME FACTORS!!!}$$

Banks


Numerical examples:

$$\Lambda_I = 8.6 \times 10^{16} \text{ GeV} \left(\frac{\alpha_{YM}}{1/25} \right)^{3/4} g^{3/4}$$

$$\Lambda_{HW} = 7.6 \times 10^{16} \text{ GeV} \left(\frac{\alpha_{YM}}{1/25} \right)^{-1/4} (4M_{GUT} V_6^{1/6})^{-3/2}$$

$$\begin{aligned} \Gamma &= \int d^4x \left\{ \frac{1}{2} m_p^2 \partial \psi \partial \psi - \Lambda^4 V(\psi) \right\} \\ &= \int d^4x \left\{ \frac{1}{2} \partial \phi \partial \phi - \Lambda^4 V(\phi / m_p) \right\} \end{aligned}$$


- The shape of the potential



zero CC min. & potential vanishes @ infinity \rightarrow intermediate max.

Inflation: constraints & predictions

- Topological inflation



δ - wall thickness in space

$$\Delta/\delta \sim \Lambda^4$$

$$H^2 \sim 1/3 \Lambda^4/m_p^2$$

Inflation $\Leftrightarrow \delta H > 1 \Leftrightarrow \Delta > m_p$

• Sufficient inflation

$$\text{Slow-roll parameters} \quad \varepsilon \sim 2 \frac{m_p^2}{4\mu} (\phi - \phi_{\max})^2, \eta \sim V''(\phi_{\max})$$

$$\mu^2 = \frac{2m_p^2}{|V''(\phi_{\max})|}$$

The “small” parameter

$$\text{Number of e-folds} \quad N(\phi, \phi_{\text{end}}) = \frac{1}{\sqrt{2}m_p} \int_{\phi}^{\phi_{\text{end}}} \frac{1}{\sqrt{\varepsilon(\phi)}} d\phi$$

$$\text{Sufficient inflation} \quad \phi_{\text{init}} - \phi_{\max} \cong \mu \text{Exp}(-[120(m_p / \mu)^2])$$

$$\text{Quantum fluct. not too large} \quad \phi_{\text{init}} - \phi_{\max} \prec \frac{H}{2\pi} \Rightarrow V''(\phi_{\max}) \prec 1/6$$

•CMB anisotropies and the string scale

$$\phi_{CMB} - \phi_{\max} = \mu \text{Exp}(-[100(m_p / \mu)^2]) \quad N(\phi_{CMB}, \phi_{\text{end}}) = 50$$

$$P_{\zeta}^{1/2} = \frac{\sqrt{3}}{2\pi} \frac{\mu \Lambda^2}{m_p^3} \text{Exp}([100(m_p / \mu)^2]) \quad P_{\zeta}^{1/2} = \frac{\sqrt{3}}{2\pi} \frac{1}{m_p^3} \frac{V^{3/2}}{V'} = \frac{15}{2} \frac{\delta\rho}{\rho}$$

$$M_S^3 / m_p = \Lambda^2 = \frac{4\pi}{\sqrt{3}} P_{\zeta}^{1/2} \frac{m_p^3}{\mu} \text{Exp}(-[100(m_p / \mu)^2])$$

$$\Lambda^2 \cong 6.5 \times 10^{16} \text{ GeV} (V''(\phi_{\max}) / 2)^{1/4} \text{Exp}(-[25V''(\phi_{\max})])$$

For consistency need $V'' \sim 1/25$

$$\varepsilon_{\text{CMB}} \propto (V'/V)^2 \sim 0$$

For our model

$$\eta_{\text{CMB}} \approx V''(\phi_{\text{max}})$$

$$n_s = 1 - 4\varepsilon_{\text{CMB}} + 2\eta_{\text{CMB}}$$

$$n_s = .92 - .08(25V''(\phi_{\text{max}}) - 1)$$

$\rightarrow \rightarrow$

$$r = 13.7 \varepsilon_{\text{CMB}}$$

$$r \approx 0$$

$$.76 < n_s < .97$$

$$1/3 < 25V'' < 3 \rightarrow$$

$$r \approx 0$$

If consistent:

$$M_S \approx 1.7 \times 10^{17} \text{ GeV} \left(\frac{P_s^{1/2}}{10^{-4}} \right)^{1/3} (1 - n_s)^{-1/6} e^{-\frac{25}{3}(1 - n_s)}$$

Summary and prospects

- Scenario: Stabilization in Central region
- Consistent cosmology:
 - scaling arguments
 - Curvature of potential needs to be “smallish”
- Predictions for CMB
- ❖ Calculate ? possible to some extent